

# Comparison of GOCE-GPS gravity fields derived by different approaches

O. Baur<sup>1</sup>, H. Bock<sup>2</sup>, P. Ditmar<sup>3</sup>, H. Hashemi Farahani<sup>3</sup>, A. Jäggi<sup>2</sup>, T. Mayer-Gürr<sup>4</sup>, T. Reubelt<sup>5</sup>, N. Zehentner<sup>4</sup>

(1) Space Research Institute, Austrian Academy of Sciences, Graz (2) Astronomical Institute, University of Bern, Switzerland (3) Delft Institute for Earth-Oriented Space research, Delft University of Technology, The Netherlands (4) Institute of Theoretical Geodesy and Satellite Geodesy, Graz University of Technology, Austria (5) Institute of Geodesy, University of Stuttgart, Germany




## Summary

Several approaches have been proposed to extract gravity field information from the GPS-derived kinematic GOCE (Gravity field and steady-state Ocean Circulation Explorer) orbits. Although there is a general consensus that, except for energy balance, these methods theoretically provide equivalent results, GOCE-GPS solutions based on real data have never been compared with each other within a consistent data processing environment so far. This contribution strives to close this gap. The gravity field solutions considered here make use of the

CMA	Celestial Mechanics Approach [1]	computed at AIUB (U Bern)
SAA	Short-Arc Approach [2]	computed at ITSG (TU Graz)
AAA	Averaged Acceleration Approach [3]	computed at DEOS (TU Delft)
PAA	Point-wise Acceleration Approach [4]	computed at GIS/IWF (U Stuttgart/Austrian Acad. of Sciences)
EBA	Energy Balance Approach [5]	computed at INAS (TU Graz)

## Formal errors; empirical errors & geoid height differences (w.r.t. ITG-Grace2010s)




## Processing details


	CMA	SAA	AAA	PAA	EBA
Orbit data	ESA SST_PKI product (reprocessed kinematic GOCE orbit)				
Variance-covariance data	ESA SST_PCV product				
Period	1.11.2009-11.1.2010 (R1)				
Spectral resolution	130				
Regularization	no				
A priori information	EGM96				
Background models	according to IERS Conventions 2003/2010				
Non-gravitational accel.	yes		no		yes
Empirical accelerations	yes			no	

## Degree-error RMS; geoid error


Orders  $\leq |0.5\pi - I|$  (inclination  $I$  in rad) omitted [6]



All orders considered




Accumulated geoid height errors




## SLR tracking residuals (obs.-comp.)

Parameterization: monthly arcs  
Estimated parameters: state vectors (1/arc), station coordinates (1/arc), drag coefficients (1/day), constant empirical accelerations (1/day), measurement biases (1/station and arc)

Lageos1 (up to degree and order 20)



c20 coefficient replaced by SLR-derived value



## Acknowledgements & References

The IWF acknowledges support by Sandro Krauss (computation of orbit residuals) and Eduard Hock (provision of the INAS solutions). The GIS thanks Matthias Roth and Matthias Weigelt for data preparation and data processing discussion, respectively.

- [1] Jäggi A, Bock H, Prange L, et al. (2011) GPS-only gravity field recovery with GOCE, CHAMP, and GRACE. Adv Space Res 47: 1020-1028
- [2] Mayer-Gürr T (2006) Gravitationsfeldbestimmung aus der Analyse kurzer Bahnbögen am Beispiel der Satellitenmissionen CHAMP und GRACE. Institut für Geodäsie und Geoinformation 9, U Bonn
- [3] Ditmar P, van Eck van der Sluijs A (2004) A technique for modeling the Earth's gravity field on the basis of satellite accelerations. J Geod 78: 12-33
- [4] Baur O, Reubelt T, Weigelt M, et al. (2012) GOCE orbit analysis: Long-wavelength gravity field determination using the acceleration approach. Adv Space Res 50: 385-396
- [5] Pail R, Bruinsma S, Migliaccio F, et al. (2011) First GOCE gravity field models derived by three different approaches. J Geod 85: 819-843
- [6] van Gelderen M, Koop R (1997) The use of degree variances in satellite gradiometry. J Geod 71: 337-343